

Mixing instructions

Proceed as follows:

1. Mix all preparations in the syringe, except NoroTec™ micronutrients.
2. Take a sample of 5 litres of spray liquid (for example at the drain plug).
3. Measure and add the appropriate amount of NoroTec micronutrients according to the table below.
4. Mix well and let stand for at least 5 minutes, preferably longer.
5. Look in the bucket to see layering, flocculation or precipitation.
6. Pour the contents into the syringe, through the strainer.
Look for precipitation.
7. If no problem is found, pour the NoroTec™ micronutrient into the syringe in the appropriate amount.
8. Treat!

Amount of fluid litres/ha	Dos NoroTec™ micronutrients litres/ha	Amount of NoroTec™ micro-nutrition in the 5-litre sample
50 litres	0,5 litres	50 ml
50 litres	1,0 litres	100 ml
50 litres	2,0 litres	200 ml
100 litres	0,5 litres	25 ml
100 litres	1,0 litres	50 ml
100 litres	2,0 litres	100 ml
150 litres	0,5 litres	17 ml
150 litres	1,0 litres	33 ml
150 litres	2,0 litres	66 ml
200 litres	0,5 litres	12,5 ml
200 litres	1,0 litres	25 ml
200 litres	2,0 litres	50 ml
250 litres	0,5 litres	10 ml
250 litres	1,0 litres	20 ml
250 litres	2,0 litres	40 ml